



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).



$\left\{ \frac{2np \Delta}{apm+bn-clp+cm}, \frac{2n \Delta}{apm+bn-clp+cm}, -\frac{2 \Delta (lp-m)}{apm+bn-clp+cm} \right\}$  are the co-ordinates of  $C$ ;

$\left\{ 0, \frac{2 \Delta (qn-1)}{b(qn-1)+cmq}, \frac{2 \Delta mq}{b(qn-1)+cmq} \right\}$  are the coordinates of  $B$ ;

$\left\{ \frac{2 \Delta (qn+1)}{a(qn+1)-qcl}, 0, -\frac{2 \Delta ql}{a(qn+1)-qcl} \right\}$  are the coordinates of  $D$ .

$a mq + \beta pmq - \gamma p(qn-1) = 0$ , is the line through  $AB$ ;

$a nql - \beta (qmn + m - lp) + \gamma n(qn+1) = 0$ , is the line through  $CD$ .

Comparing, we get  $q=0$  or  $q=- (1/n)$ ,  $p=m/l$  or  $p=0$ . There are no positive values for  $q$  when  $p$  is positive. When  $p$  is positive,  $q=0$ , etc.

Whatever relations we establish we cannot find  $p$  and  $q$  both real or both positive.

## CALCULUS.

286. Proposed by R. D. CARMICHAEL, Princeton University.

Solve the differential equation

$$\begin{aligned} & [a_0 x^3 + a_1 x^2 y + a_2 x y^2 + (a_0 - a_1 + a_2) y^3 \\ & \quad + a_3 x^2 + a_4 x y + a_5 y^2 + a_6 x + a_7 y + a_8] dx \\ & + [a_0 y^3 + a_1 x y^2 + a_2 x^2 y + (a_0 - a_1 + a_2) x^3 \\ & \quad + a_3 y^2 + a_4 x y + a_5 x^2 + a_6 y + a_7 x + a_8] dy = 0. \end{aligned}$$

No solution of this problem has been received.

287. Proposed by C. N. SCHMALL, 604 East 5th Street, New York City.

An object  $P$ , being placed beyond the principal focus  $F$  of a convex lense, determine its position when its distance  $PQ$ , from its image  $Q$ , is a minimum.

Solution by G. B. M. ZERR, A. M., Ph. D., Philadelphia, Pa.

Let  $u$ =distance of object from lense,  $v$ =distance of image from lense,  $t$ =thickness of lense, and  $r, s$ =the radii of the first and second surface.

Then  $\frac{1}{\frac{1}{v} + \frac{\mu-1}{s}} + \frac{1}{\frac{1}{u} - \frac{\mu-1}{r}} = \frac{t}{\mu}$ , where  $\mu$ =index of refraction, or

$$\mu \left( \frac{1}{u} + \frac{1}{v} \right) = \left( \frac{1}{r} - \frac{1}{s} \right) \mu (\mu - 1) + t \left( \frac{1}{v} + \frac{\mu-1}{s} \right) \left( \frac{1}{u} - \frac{\mu-1}{r} \right) \dots (1).$$